

SMMCI -Case series FINAL

29.4.2021.docx

By Eliyahu Mass

Abstract:

¹⁹ Solitary median maxillary central incisor (SMMCI) is a rare oddity, which may be associated with a complex of multiple developmental abnormalities. Sometimes, it is related to background systemic features, but may also be a separate localized anatomical aberration. Its occurrence has been estimated in 1:50,000 live births. SMMCIs were described together with cleft lip and palate (CLP), Velo Cardio Facial Syndrome (VCFS), exaggerated midline maxillary torus, distorted soft palate, and absence of the uvula and the midline labial frenum. Associated with systemic conditions, it has also been described with hypotelorism, hypothalamic-pituitary disorders, coloboma, ectodermal dysplasia and blindness secondary to microphthalmia. It occurs ³³ in both primary and permanent dentitions of the same child. Nasal malformation like flat nose with a single nostril, choanal atresia, mid-nasal stenosis, or ²² congenital nasal pyriform aperture stenosis (CNPAS) may also be associated with the disorder. In severe cases, it may be associated with holoprosencephaly (HPE) and cyclopia associated with Ito hypomelanosis. We present occasional sporadic cases of otherwise normal children.

21

Introduction:

Solitary median maxillary central incisor (SMMCI) as an isolated occurrence and SMMCI as part of a syndrome are rare dental anomalies. (1) The incidence has been estimated as 1:50,000 live births. (2) These anomalies are caused by heterozygous mutation in the Sonic Hedgehog Gene (SHH; 600725) on chromosome 7q36. (3)

The awareness of these anomalies goes back to 1958, where Scott reported "Absence of upper central incisor". (3) Since then, the scientific literature regarding SMMCI is characterized by single or sporadic case reports, as well as efforts to investigate the etiology and delineate associated developmental and genetic aberrations.

Short stature and deficient growth hormone were described in patients with solitary maxillary central incisors in the primary and consecutive permanent dentitions. In some cases with this feature, growth hormones were normal. No similar or associated oral and dental aberrations were found in families of those patients. Superoincisivodontic dwarfism, also known as 'monosuperoincisivodontic dwarfism' was used then to describe such phenomena. (4,5) Choanal atresia/midnasal stenosis and holoprosencephaly (HPE) was added to the short stature to a series of 21 consecutive cases of SMMCI syndrome in 1997. (2)

The holoprosencephaly spectrum (HPE): HPE stands for a teratologic series of varying severity, expressed by median malformations of the face and brain. There are a variety of chromosomal causes, yet most cases are sporadic and of unknown etiology. Less severe cases of Autosomal Dominant (AD) HPE

may present with mild manifestations of hypotelorism, anosmia, hyposmia (reduced ability to smell), microcephaly, mental deficiency, or midface hypoplasia, and SMMCI. Ocular colobomas may also be included.

³ HPE is a specific malformation complex resulting from incorrect cleavage of the embryonic prosencephalon, (forebrain) with concurrent facial anomalies (midline facial structures) ranging from cyclopia to mild forms such as hypotelorism and SMMCI, with other clinical manifestations such as anophthalmia to mild hypotelorism, coloboma, hyposmia, dysplasia of olfactory bulbs and optic nerves. ⁽⁶⁾

The inheritance of HPE is Autosomal Dominant – the parents might be with minor manifestations such as single central incisor, hypotelorism, hyposmia, but offspring may be with severe manifestations, which emphasizes the importance of genetic counseling.

¹ SMMCI syndrome, previously considered a simple midline defect of the dental lamina is "a possible predictor of HPEs of varying degrees in the proband, in members of the proband's family, and in the family's descendants". ⁽²⁾

³² Most cases of HPE occur sporadically, yet occasionally a single upper central incisor associated with HPE may be found in close relatives (kindred). These findings "may represent evidence for a less severe form of HPE that may be transmitted in an AD fashion". ⁽⁷⁾

²³ Midline craniofacial malformations in association with hypothalamic-pituitary disorders has been observed in a girl with true precocious puberty associated with a SMMCI in both primary and permanent dentitions, and a hypothalamic hamartoma. ⁽⁸⁾

Genetics: Dolan et al (1981) described a 28-month-old boy, and Aughton et al. (1991) reported a 7 years old girl with SMMCI. (9,10) Through genetic examination, both were found to have (18p) deletion syndrome. Due to the rarity of reporting of the anomaly at that time (1981-1991), SMMCI were evaluated as fused maxillary central incisors. It is associated with some degree of HPE in some cases with SMMCI in the mild manifestation of the HPE. (10)

Occasionally, SMMCI with semilobar HPE, a median cleft lip, flat nose with a single nostril, hypotelorism, and normal chromosomes, may be an incidental mutation in otherwise healthy close relative, with no apparent dental or facial anomalies. Indeed, the presence of SMMCI in a newborn baby may be significant in genetic counseling as an indicator of potential HPE in the next generation, even if other relatives are apparently normal. The reason behind this suggestion is that the penetrance and expressivity of the AD form of HPE vary widely. (11)

Winter et al. (1988), (12) and Buntinx and Baraitser (1989) (13) described a SMMCI in patients with a form of ectodermal dysplasia (ED). Although this is an unusual manifestation of ED, it should be considered as significant in possible gene carriers. In the family presented there, the inheritance pattern is likely to be autosomal recessive (AR). (12,13) Artman and Boyden (1990) (14) described a 5 years 1 month old girl with 'slow growth', periodic severe headaches, intermittent nasal congestion, and near complete blindness secondary to microphthalmia. Her height and weight were both well below 3 SD and at the 50th centile for an 18 month and 9 months respectively. Her bone age was of 2 years 5 months and CT scan showed a small sella turcica with a hypoplastic pituitary gland. A small bony ridge was noted in the

roof of the hard palate. The growth hormone was below that of normal controls. The girl has been defined as having isolated growth hormone deficiency, microphthalmia and isolated growth hormone deficiency in association with SMMCI. (14)

The association of SMMCI with nasal obstruction was recognized. (15)
Congenital nasal pyriform aperture stenosis (CNPAS) was described by Brown in 1989 (16) as a cause of nasal airway obstruction in the newborn, but there was no mention of SMMCI. (16)

SMMCI is a non-specific sign, it may occur as an isolated abnormality or as microform of HPE with autosomal dominant transmission. (15) Those patients noted to suffer from congenital nasal pyriform aperture stenosis (CNPAS) must be observed for eruption of a SMMCI, which has been described as "megaincisor" together with maxillary bony overgrowth. These patients must be assumed to have a microform of HPE, which necessitates chromosomal analysis, evaluation of relatives, counseling of parents, CT scan to check for CNS malformations, and assessment of the hypothalamic-pituitary-thyroid-adrenal axis. Indeed, CNPAS may suggest a midfacial dysostosis with associated endocrine and central nervous system abnormalities, more than an isolated congenital abnormality of the airway.

SMMCI may occur as an isolated finding or in association with other systemic abnormalities like short stature, pituitary insufficiency, microcephaly, choanal atresia, midnasal stenosis, and CNPAS. It can also be a feature of recognized syndromes with specific chromosomal abnormalities or with no chromosomal abnormality. (17)

¹ Masuno et al. (1990) ⁽¹⁸⁾ described 2 unrelated cases of SMMCI with 7q terminal deletion. ⁸ They had mental retardation, microcephaly, hypotelorism, short stature, and normal levels of plasma growth hormone. One patient had bilateral caudal ectopic kidneys, double renal pelvises, and dilated ureters. The other had bilateral hydronephrosis. The authors suggested that 7q terminal deletion is one of the causes of ¹ SMMCI. ⁽¹⁸⁾

Nanni et al. (2001) ⁽¹⁷⁾ performed a molecular study of 13 patients with SMMCI ¹ who did not have HPE. They studied 2 genes, SHH (600725) and SIX3 (603714), in which mutations had been reported in patients showing SMMCI as part of the HPE spectrum. They found a new missense mutation in SHH (I111F; 600725.0014) which may be specific for the SMMCI phenotype since it had not been found in patients with HPE or in normal controls. ⁽¹⁷⁾

Marini et al. (2003) ⁽¹⁹⁾ studied a family, previously described by Camera et al. (1992), ⁽²⁰⁾ in which the mother presented with a SMMCI ¹ and mild hypotelorism and her daughter and 2 fetuses were diagnosed with HPE. Sequencing of DNA in this family identified a nonsense mutation in the SHH gene. ^(19,20) In this paper we present cases, otherwise healthy children with this phenomenon and one suspected with systemic aberrations.

About 30 years ago (1991), one of us (EM) came across a boy with SMMCI phenomenon, which proved to be unfamiliar to most peer colleagues across the country (Figures 1,2). ⁽²¹⁾ By now, this case, has 2 otherwise healthy children. Like his mother then, he refuses systemic or genetic examinations. Other few cases were provided by sporadic practitioners. Only one additional child, with obvious features, suggesting a syndrome (case EM), has been examined for dental treatment, but the mother had refused further

comprehensive examinations, which prevented detailed information. Hence, the purpose of this review and presentation of further small case series is to draw attention to the rarity of this phenomenon, sometimes referred to as a syndrome, or a part of a genetic feature with, or without variable associated anomalies.

Case series:

First two cases (Figures 1,2) were presented in details in Mass and Sarnat 1991 (21).

Case no 3:

KS – A 7 years old boy was referred by a general practitioner for consultation, due to "Delayed dental development and a peculiar phenomenon in the maxillary incisal area". Other than the characteristic appearance in the primary and the following permanent SMMCI and exaggerated midline palatal torus in a maxillary cohesion, medical and dental examinations were inconclusive (Figure 3).

Case no 4:

EM - A 5 years old girl was referred for dental examination and treatment. The examination revealed mid nasal bulge; mild hypotelorism (in contrast, her brother had mild hypertelorism); atypical soft palate and lack of uvula; lack of anterior frenum; prominent mid palatal torus, without incisive papilla (Figure 4).

Upon suggestion to perform further systematic examination, the mother refused, leaving the clinic, claiming that she brought the child for "fillings".

Summary and Conclusions:

Background: SMMCI and SMMCI syndrome is a rare phenomenon presented in the scientific literature sporadically. Due to its rarity, it was felt that any additional reports may contribute to contemporary awareness of the importance of comprehensive case examinations.

Aim:

⁷ This article describes a case series of additional cases with common features, involving a solitary median maxillary central incisor. The patients were seen by different practicing dentists in Israel. Of these cases, none went through genetic and/or systemic examination, due to unawareness of its genetic and systemic implications, which otherwise were not noticed, and by refusal of one of the parents for further comprehensive examinations. Yet, 2 features, which were not mentioned in previous reports, are tendency of the root apex of the permanent SMMCI to split up and the lack of visible incisive papilla. Indeed, ⁷ it is now recognized as a possible predictor of holoprosencephalies of varying degrees in members of the proband's family, and in the family's descendants. Hence, it is hoped that this literature review and the presented cases ⁷ will help to clarify understanding of solitary median maxillary central incisor as a sporadic entity or as a syndrome, and prompt pediatric dentists to conduct thorough tests upon its detection.

Conclusions:

In a world of advanced technology and communication, any additional information may provide potential clinical significance and expand understanding, which may improve professional approach.

List of figures:

Figure 1. TA - (21) Anterior periapical radiograph; Note splitting of the pulp canal close to the apex (type IV according to Weine's classification. (22)

Figure 2. RM - Anterior periapical radiograph during orthodontic treatment. (21) Note splitting of the pulp canal close to the apex (type IV according to Weine's classification. (22)

Figures 3. – Facial and dental clinical pictures and anterior periapical radiograph of case KS. Note the exaggerated midline palatal torus, lack of anterior labial frenum and incisive papilla.

Figure 4. – Facial and dental clinical pictures and anterior periapical radiograph of case EM. Note the exaggerated midline palatal torus, lack of anterior labial frenum, incisive papilla, posterior uvula with flabby soft palate.

REFERENCES:

1. ¹⁰ Garavelli L, Zanacca C, Caselli G, Banchini G, Dubourg C, David V, et al. Solitary median maxillary central incisor syndrome: clinical case with a novel mutation of sonic hedgehog. *Am J Med Genet* 2004; 127A: 93-95.
2. ² Hall RK, Bankier A, Aldred MJ, Kan K, Lucas JO, Perks AGB. Solitary median maxillary central incisor, short stature, choanal atresia/midnasal stenosis (SMMCI) syndrome. *Oral Surg Oral Med Oral Path Oral Radiol Endod* 1997; 84: 651-662.
3. ³ Scott DC. Absence of upper central incisor. *Br Dent J* 1958; 104: 247-248. Online Mendelian In Men (OMIM # 147250).
4. ²⁶ ¹ Rappaport EB, Ulstrom RA, Gorlin RJ. Monosupero-centroincisivodontic dwarfism. *Birth Defects Orig Art Ser* 1976; XII(5) :243-245.
5. ⁵ Rappaport EB, Ulstrom RA, Gorlin RJ, Lucky AW, Colle E, Miser J. Solitary maxillary central incisor and short stature. *J Pediat* 1977; 91: 924-928.
6. ² Liberfarb RM, Abdo OP, Pruett RC. Ocular coloboma associated with a solitary maxillary central incisor and growth failure: manifestations of holoprosencephaly. *Ann Ophthal* 1987; 19: 226-227.
7. ¹⁷ ⁷ Berry SA, Pierpont ME, Gorlin RJ. Single central incisor in familial holoprosencephaly. 1984; *J Pediat* 104: 877-880.
8. ⁶ ⁸ Winter WE, Rosenbloom AL, MacLean NK, Mickle PJ. Solitary, central, maxillary incisor associated with precocious puberty and hypothalamic hamartoma. *J Pediat* 1982; 101: 965-967.
9. ⁵ ⁹ Dolan LM, Willson K, Wilson WG. 18p- syndrome with a single central maxillary incisor. *J Med Genet* 1981; 18: 396-398.

10. Aughton DJ, AlSaadi AA, Transue DJ. Single maxillary central incisor in a girl with del(18p) syndrome. *J Med Genet* 1991; 28: 530-532.
11. Hattori H, Okuno T, Momoi T, Kataoka K, Mikawa H, Shiota K. Single central maxillary incisor and holoprosencephaly. *Am J Med Genet* 1987; 28: 483-487 [PubMed: 3425622, related citations]
12. Winter RM, MacDermot KD, Hill FJ. Sparse hair, short stature, hypoplastic thumbs, single upper central incisor and abnormal skin pigmentation: a possible 'new' form of ectodermal dysplasia. *Am J Med Genet* 1988; 29: 209-216.
13. Buntinx I, Baraitser MA. Single maxillary incisor as a manifestation of an ectodermal dysplasia. *J Med Genet* 1989; 26: 648-651.
14. Artman HG, Boyden E. Microphthalmia with single central incisor and hypopituitarism. *J Med Genet* 1990; 27: 192-193.
15. Arlis H, Ward RF. Congenital nasal pyriform aperture stenosis: isolated abnormality vs developmental field defect. *Arch Otolaryng Head Neck Surg.* 1992; 118: 989-991.
16. Brown OE, Myer III CM, Manning SC. Congenital nasal pyriform aperture stenosis. *Laryngoscope* 1989; 99: 86-91.
17. Nanni L, Ming JE, Du Y, Hall RK, Aldred M, Bankier A, Muenke M. SHH mutation is associated with solitary median maxillary central incisor: a study of 13 patients and review of the literature. *Am J Med Genet* 2001; 102: 1-10.
18. Masuno M, Fukushima Y, Sugio Y, Ikeda M, Kuroki Y. Two unrelated cases of single maxillary central incisor with 7q terminal deletion. *Jpn J Hum Genet* 1990; 35: 311-317.
19. Marini M, Cusano R, De Biasio P, Caroli F, Lerone M, Silengo M, Ravazzolo R, Seri M, Camera G. Previously undescribed nonsense mutation in

SHH caused autosomal dominant holoprosencephaly with wide intrafamilial variability. *Am J Med Genet* 2003; 117A: 112-115.

¹
20. Camera G, Bovone S, Zucchinetti P, Pozzolo S, Giunta E. Incisivo mascellare centrale unico e eolprosencefalia. *Pathologica* 1992; 84: 425-428.

²
21. Mass E, Sarnar H. Single maxillary central incisor in the midline. *J Dent Child*; Sep. Oct. 1991, 413-416.

¹²
22. Weine FS, Healey HJ, Gerskin H, Evanson L. Canal configuration in the mesibuccal root of maxillary first molar and its endodontic significance. *Oral Surge Oral Med Oral Pathol* 1969; 28: 419-425.

46%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|---|---|-----------------|
| 1 | uwm.edu
Internet | 284 words — 11% |
| 2 | www.ojrd.com
Internet | 203 words — 8% |
| 3 | Arlis, H., and R. F. Ward. "Congenital Nasal Pyriform Aperture Stenosis: Isolated Abnormality vs Developmental Field Defect", Archives of Otolaryngology - Head and Neck Surgery, 1992.
Crossref | 94 words — 4% |
| 4 | H G Artman, E Boyden. "Microphthalmia with single central incisor and hypopituitarism.", Journal of Medical Genetics, 1990
Crossref | 71 words — 3% |
| 5 | Serap Turan, Nurcin Saka, Ilter Guney, Abdullah Bereket. "A Patient with Hypopituitarism and Isochromosome 18q Mosaicism", Hormone Research in Paediatrics, 2005
Crossref | 70 words — 3% |
| 6 | jmg.bmj.com
Internet | 52 words — 2% |
| 7 | lrd.yahooapis.com
Internet | 47 words — 2% |

-
- 8 Mitsuo Masuno. "Two unrelated cases of single maxillary central incisor with 7q terminal deletion", The Japanese Journal of Human Genetics, 12/1990
Crossref 41 words — 2%
-
- 9 Marion Devambez, Alexis Delattre, Pierre Fayoux. "Congenital Nasal Pyriform Aperture Stenosis: Diagnosis and Management", The Cleft Palate-Craniofacial Journal, 2009
Crossref 38 words — 1%
-
- 10 www.labome.org
Internet 35 words — 1%
-
- 11 Raquel Garcia Rodriguez, Loida Garcia Cruz, Yeray Novoa Medina, Raquel Garcia Delgado et al. "The solitary median maxillary central incisor (SMMCI) syndrome: Associations, prenatal diagnosis, and outcomes", Prenatal Diagnosis, 2019
Crossref 30 words — 1%
-
- 12 app.trdizin.gov.tr
Internet 29 words — 1%
-
- 13 Luisa Nanni, Jeffrey E. Ming, Yangzhu Du, Roger K. Hall, Michael Aldred, Agnes Bankier, Maximilian Muenke. "SHH mutation is associated with solitary median maxillary central incisor: A study of 13 patients and review of the literature", American Journal of Medical Genetics, 2001
Crossref 24 words — 1%
-
- 14 Micky Osovsky, Anat Aizer-Danon, Gadi Horev, Lea Sirota. "Congenital pyriform aperture stenosis", Pediatric Radiology, 2006
Crossref 21 words — 1%

-
- 15 Haruo Hattori. "Single central maxillary incisor and holoprosencephaly", American Journal of Medical Genetics, 10/1987 17 words — 1%
Crossref
-
- 16 mafiadoc.com 17 words — 1%
Internet
-
- 17 oatext.com 16 words — 1%
Internet
-
- 18 Haruo Hattori, Takehiko Okuno, Toru Momoi, Kenkichi Kataoka, Haruki Mikawa, Kohei Shiota, John M. Opitz, James F. Reynolds. "Single central maxillary incisor and holoprosencephaly", American Journal of Medical Genetics, 1987 16 words — 1%
Crossref
-
- 19 apjai.digitaljournals.org 12 words — < 1%
Internet
-
- 20 www.questia.com 10 words — < 1%
Internet
-
- 21 Van Dijk, F.S.. "Solitary median maxillary central incisor and congenital nasal pyriform aperture stenosis combined with asymmetric crying facies and postaxial lower limb reduction defects: A unique combination of features", European Journal of Medical Genetics, 201105/06 10 words — < 1%
Crossref
-
- 22 Guillaume Captier. "Anatomical Consideration of the Congenital Nasal Pyriform Aperture Stenosis: Localized Dysostosis Without Interorbital Hypoplasia", Journal of Craniofacial Surgery, 05/2004 10 words — < 1%
Crossref

23 William E. Winter, Arlan L. Rosenbloom, Noel K. Maclaren, J. Parker Mickle. "Solitary central maxillary incisor associated with precocious puberty and hypothalamic hamartoma", The Journal of Pediatrics, 1982
Crossref 9 words — < 1%

24 www.aapd.org
Internet 9 words — < 1%

25 Snehlata Oberoi. "Velocardiofacial syndrome with single central incisor", American Journal of Medical Genetics Part A, 01/15/2005
Crossref 8 words — < 1%

26 I Buntinx, M Baraitser. "A single maxillary incisor as a manifestation of an ectodermal dysplasia.", Journal of Medical Genetics, 1989
Crossref 7 words — < 1%

27 "The Pigmentary System", Wiley, 2006
Crossref 7 words — < 1%

28 Robin M. Winter, Michael Baraitser. "Chapter 1 A catalogue of multiple congenital anomaly syndromes", Springer Science and Business Media LLC, 1991
Crossref 7 words — < 1%

29 Brian C. Harsha. "Solitary central maxillary incisor associated with precocious puberty and hypothalamic hamartoma", Journal of Oral and Maxillofacial Surgery, 1983
Crossref 7 words — < 1%

30 Francisco Wanderley Garcia de Paula e Silva. "Solitary median maxillary central incisor in association with Goldenhar's syndrome: a case report", Special Care in Dentistry, 5/2007
7 words — < 1%

31 "Atlas of Genetic Diagnosis and Counseling", Springer Science and Business Media LLC, 2006 7 words — < 1%
Crossref

32 Susan A. Berry, Mary Ella Pierpont, Robert J. Gorlin. "Single central incisor in familial holoprosencephaly", The Journal of Pediatrics, 1984 6 words — < 1%
Crossref

33 Roger K Hall. "Solitary median maxillary central incisor (SMMCI) syndrome", Orphanet Journal of Rare Diseases, 2006 6 words — < 1%
Crossref

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY OFF

EXCLUDE MATCHES OFF